

Cavity QED and Quantum Computation in the Strong Coupling Regime II : Two Atoms Case

Kazuyuki FUJII *

Department of Mathematical Sciences

Yokohama City University

Yokohama, 236-0027

Japan

Abstract

In this paper we study the behavior of (laser-cooled) m -atoms trapped in a cavity interacting with a photon $\cdots$ Cavity QED $\cdots$ and attempt to solve the Schrödinger equation of this model in the strong coupling regime. In the case of $m = 2$ we construct Bell-Schrödinger cat states (in our terminology) and obtain with such bases some unitary transformations by making use of the rotating wave approximation under new resonance conditions containing the Bessel functions, which will be applied to construct important quantum logic gates in Quantum Computation. Moreover we propose in the case of $m = 3$ a crucial problem to solve on Quantum Computation.

This paper is a continuation of [15], [16]. We consider a unified model of the interaction of the two-level atoms and both the single radiation mode and external field (periodic usually) in a cavity. We deal with the external field as a classical one. As a general

*E-mail address : fujii@yokohama-cu.ac.jp

introduction to this topic in Quantum Optics see [1], [2], [3]. Our model is deeply related to the one of trapped ions in a cavity with the photon interaction (Cavity QED).

In our model we are especially interested in the strong coupling regime, [8], [13], [14]. One of motivations is a recent very interesting experiment, [11]. See [3] and [9] as a general introduction.

In [8] and [13] we have treated the strong coupling regime of the interaction model of the two-level atom and the single radiation mode, and have given some explicit solutions under the resonance conditions and rotating wave approximations.

On the other hand we want to add some external field (like Laser one) to the above model which will make the model more realistic (for example in Quantum Computation). Therefore we propose the unified model.

We would like to solve our model in the strong coupling regime. Especially we want to show the existence of Rabi oscillations in this regime because the real purpose of a series of study ([13], [14], [15], [16]) is an application to Quantum Computation (see [12] as a brief introduction to it).

Since in the previous paper [16] we have treated one-atom case, so we deal with two-atoms case in this paper. To solve the Schrödinger equation in this regime we construct (in our terminology) Bell-Schrödinger cat states, and make use of rotating wave approximation under new resonance conditions containing the Bessel functions and etc, and obtain unitary transformations which are necessary to perform quantum logic gates.

Our solutions might give a new insight into Quantum Optics or Condensed Matter Physics as well as Quantum Computation.

We here make some preparations for the latter. Let $\{\sigma_1, \sigma_2, \sigma_3\}$ be Pauli matrices and $\mathbf{1}_2$ a unit matrix :

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathbf{1}_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad (1)$$


and $\sigma_+ = (1/2)(\sigma_1 + i\sigma_2)$, $\sigma_- = (1/2)(\sigma_1 - i\sigma_2)$. Let W be the Walsh–Hadamard matrix

$$W = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} = W^{-1}, \quad (2)$$

then we can diagonalize σ_1 as $\sigma_1 = W\sigma_3W^{-1} = W\sigma_3W$ by making use of this W . The eigenvalues of σ_1 is $\{1, -1\}$ with eigenvectors

$$|1\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad |-1\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \end{pmatrix} \implies |\lambda\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ \lambda \end{pmatrix}. \quad (3)$$

We have treated one atom with two–level, so we would like to generalize the method developped in [13], [15] to m atoms with two–level interacting both the single radiation mode and (classical) external fields like (m atoms trapped in a cavity)



Then the unified Hamiltonian can be given as

$$H_L = \omega \mathbf{1}_M \otimes L_3 + g_1 \sum_{j=1}^m \sigma_1^{(j)} \otimes (L_+ + L_-) + \frac{\Delta}{2} \sum_{j=1}^m \sigma_3^{(j)} \otimes \mathbf{1}_L + g_2 \sum_{j=1}^m \cos(\omega_j t + \phi_j) \sigma_1^{(j)} \otimes \mathbf{1}_L, \quad (4)$$

where $M = 2^m$ and $\sigma_k^{(j)}$ ($k = 1, 3$) is

$$\sigma_k^{(j)} = \mathbf{1}_2 \otimes \cdots \otimes \mathbf{1}_2 \otimes \sigma_k \otimes \mathbf{1}_2 \otimes \cdots \otimes \mathbf{1}_2 \quad (j - \text{position}),$$

see [15]. Here we are treating the following three cases at the same time : The notation $\{L_+, L_-, L_3\}$ means respectively

$$\{L_+, L_-, L_3\} = \begin{cases} \text{(N)} & \{a^\dagger, a, N\}, \\ \text{(K)} & \{K_+, K_-, K_3\}, \\ \text{(J)} & \{J_+, J_-, J_3\}, \end{cases} \quad (5)$$

and $\{a^\dagger, a, N\}$ is the generators of Heisenberg algebra, $\{K_+, K_-, K_3\}$ and $\{J_+, J_-, J_3\}$ are a set of generators of unitary representations of Lie algebras $su(1, 1)$ and $su(2)$. They are usually constructed by making use of two harmonic oscillators (two-photons) $\{a_1, a_1^\dagger\}$, $\{a_2, a_2^\dagger\}$ as

$$\begin{aligned} su(1, 1) : \quad K_+ &= a_1^\dagger a_2^\dagger, \quad K_- = a_2 a_1, \quad K_3 = \frac{1}{2} (a_1^\dagger a_1 + a_2^\dagger a_2 + 1), \\ su(2) : \quad J_+ &= a_1^\dagger a_2, \quad J_- = a_2^\dagger a_1, \quad J_3 = \frac{1}{2} (a_1^\dagger a_1 - a_2^\dagger a_2). \end{aligned}$$

See for example [13], [17] in detail.

We are interested in the strong coupling regime ($g_1 \gg \Delta$), so we solve the Hamiltonian as follows (see [15]). Let us transform (4) into

$$\begin{aligned} H_L &= \omega \mathbf{1}_M \otimes L_3 + \sum_{j=1}^m \sigma_1^{(j)} \otimes \{g_1(L_+ + L_-) + g_2 \cos(\omega_j t + \phi_j) \mathbf{1}_L\} + \frac{\Delta}{2} \sum_{j=1}^m \sigma_3^{(j)} \otimes \mathbf{1}_L \\ &\equiv H_0 + \frac{\Delta}{2} \sum_{j=1}^m \sigma_3^{(j)} \otimes \mathbf{1}_L, \end{aligned} \tag{6}$$

where we have written H_0 instead of H_{L0} for simplicity. First we diagonalize H_0 . For that we set

$$\mathbf{W} = W \otimes W \otimes \cdots \otimes W \quad \in \quad U(M)$$

for W in (2). Then it is not difficult to see (see [13], [15])

$$\begin{aligned} H_0 &= (\mathbf{W} \otimes \mathbf{1}_L) \left[\mathbf{1}_M \otimes \omega L_3 + \sum_{j=1}^m \sigma_3^{(j)} \otimes \{g_1(L_+ + L_-) + g_2 \cos(\omega_j t + \phi_j) \mathbf{1}_L\} \right] (\mathbf{W}^{-1} \otimes \mathbf{1}_L) \\ &= (\mathbf{W} \otimes \mathbf{1}_L) \left[\mathbf{1}_M \otimes \omega L_3 + g_1 \sum_{j=1}^m \sigma_3^{(j)} \otimes (L_+ + L_-) + g_2 \sum_{j=1}^m \sigma_3^{(j)} \otimes \cos(\omega_j t + \phi_j) \mathbf{1}_L \right] (\mathbf{W}^{-1} \otimes \mathbf{1}_L) \\ &= \sum_{\lambda} \left(|\lambda\rangle \otimes e^{-\frac{\Lambda x}{2}(L_+ - L_-)} \right) \left\{ \Omega L_3 + g_2 \sum_{j=1}^m \lambda_j \cos(\omega_j t + \phi_j) \mathbf{1}_L \right\} \left(\langle \lambda| \otimes e^{\frac{\Lambda x}{2}(L_+ - L_-)} \right), \end{aligned} \tag{7}$$

where we have used the following

Key Formulas

$$\begin{aligned} (N) \quad \omega a^\dagger a + g_1 \Lambda (a^\dagger + a) &= \Omega e^{-\frac{\Lambda x}{2}(a^\dagger - a)} \left(N - \frac{(g_1 \Lambda)^2}{\omega^2} \right) e^{\frac{\Lambda x}{2}(a^\dagger - a)} \\ \text{where} \quad \Omega &= \omega, \quad x = 2g_1/\omega, \end{aligned} \tag{8}$$

$$(K) \quad \omega K_3 + g_1 \Lambda (K_+ + K_-) = \Omega e^{-\frac{\Lambda x}{2}(K_+ - K_-)} K_3 e^{\frac{\Lambda x}{2}(K_+ - K_-)}$$

$$\text{where} \quad \Omega = \omega \sqrt{1 - (2g_1 \Lambda / \omega)^2}, \quad x = (1/\Lambda) \tanh^{-1}(2g_1 \Lambda / \omega), \quad (9)$$

$$(J) \quad \omega J_3 + g_1 \Lambda (J_+ + J_-) = \Omega e^{-\frac{\Lambda x}{2}(J_+ - J_-)} J_3 e^{\frac{\Lambda x}{2}(J_+ - J_-)}$$

$$\text{where} \quad \Omega = \omega \sqrt{1 + (2g_1 \Lambda / \omega)^2}, \quad x = (1/\Lambda) \tanh^{-1}(2g_1 \Lambda / \omega), \quad (10)$$

and have used concise notations $\lambda = (\lambda_1, \dots, \lambda_{j-1}, \lambda_j, \lambda_{j+1}, \dots, \lambda_m)$ and

$$\sum_{\lambda} = \sum_{\lambda_1=\pm 1} \sum_{\lambda_2=\pm 1} \cdots \sum_{\lambda_m=\pm 1} \quad ; \quad |\lambda\rangle = |\lambda_1\rangle \otimes |\lambda_2\rangle \otimes \cdots \otimes |\lambda_m\rangle \quad ; \quad \Lambda \equiv \Lambda(\lambda) = \sum_{j=1}^m \lambda_j$$

for $\lambda_j = \pm 1$. We leave the proof of Key Formulas to the readers.

That is, we could diagonalize the Hamiltonian H_0 . Its eigenvalues $\{E_n(t)\}$ and eigenvectors $\{|\{\lambda, n\}\rangle\}$ are given respectively

$$(E_n(t), |\{\lambda, n\}\rangle) = \begin{cases} (N) & \Omega(-\frac{g_1^2}{\omega^2} \Lambda^2 + n) + g_2 \sum_{j=1}^m \lambda_j \cos(\omega_j t + \phi_j), \quad |\lambda\rangle \otimes e^{-\frac{\Lambda x}{2}(a^\dagger - a)} |n\rangle, \\ (K) & \Omega(K + n) + g_2 \sum_{j=1}^m \lambda_j \cos(\omega_j t + \phi_j), \quad |\lambda\rangle \otimes e^{-\frac{\Lambda x}{2}(K_+ - K_-)} |K, n\rangle, \\ (J) & \Omega(-J + n) + g_2 \sum_{j=1}^m \lambda_j \cos(\omega_j t + \phi_j), \quad |\lambda\rangle \otimes e^{-\frac{\Lambda x}{2}(J_+ - J_-)} |J, n\rangle \end{cases} \quad (11)$$

for $n \in \mathbf{N} \cup \{0\}$, where $E_n(t) \equiv E_n(\lambda) + g_2 \sum_{j=1}^m \lambda_j \cos(\omega_j t + \phi_j)$. Then H_0 above can be written as

$$\begin{aligned} H_0 &= \sum_{\lambda} \sum_n E_n(t) |\{\lambda, n\}\rangle \langle \{\lambda, n\}| \\ &= \sum_{\lambda} \sum_n \left\{ E_n(\lambda) + g_2 \sum_{j=1}^m \lambda_j \cos(\omega_j t + \phi_j) \right\} |\{\lambda, n\}\rangle \langle \{\lambda, n\}|. \end{aligned}$$

Up to this stage we could treat $L = (N), (K), (J)$ at the same time. However in the following it is very difficult to treat them at the same time, so we shall focus our attention to the simplest case $L = (N)$ ¹. We leave the remaining cases as a future (challenging) task.

Next we would like to solve the following Schrödinger equation :

$$i \frac{d}{dt} \Psi = H_L \Psi = \left(H_0 + \frac{\Delta}{2} \sum_{j=1}^m \sigma_3^{(j)} \otimes \mathbf{1}_L \right) \Psi. \quad (12)$$

¹The main difficulty for $L = (K)$ and (J) is the λ dependence of x , see (9) and (10).

To solve this equation we appeal to the method of constant variation. First let us solve $i\frac{d}{dt}\Psi = H_0\Psi$, which general solution is given by $\Psi(t) = U_0(t)\Psi_0$, where Ψ_0 is a constant state and

$$U_0(t) = \sum_{\lambda} \sum_n e^{-i\left\{tE_n(\lambda) + \sum_{j=1}^m \lambda_j (g_2/\omega_j) \sin(\omega_j t + \phi_j)\right\}} |\{\lambda, n\}\rangle \langle \{\lambda, n\}|. \quad (13)$$

The method of constant variation goes as follows. Changing like $\Psi_0 \longrightarrow \Psi_0(t)$, we have

$$i\frac{d}{dt}\Psi_0 = \frac{\Delta}{2} U_0^\dagger \left(\sum_{j=1}^m \sigma_3^{(j)} \otimes \mathbf{1}_L \right) U_0 \Psi_0 \equiv \frac{\Delta}{2} H_F \Psi_0. \quad (14)$$

Then making use of the fact $\langle \lambda_j | \sigma_3 = \langle -\lambda_j |$ and after long calculations, H_F becomes

$$H_F = \sum_{\lambda} \sum_{n, n'} \sum_{j=1}^m e^{i\left\{t\omega(n-n') + t\omega x^2(1-\lambda_j\Lambda(\lambda)) + 2\lambda_j(g_2/\omega_j) \sin(\omega_j t + \phi_j)\right\}} \times \\ \langle n | e^{\lambda_j x(a^\dagger - a)} | n' \rangle |\{\lambda, n\}\rangle \langle \{\lambda, n'\}_{(j)}|, \quad (15)$$

where we have used the compact notations $\lambda_{(j)} = (\lambda_1, \dots, \lambda_{j-1}, -\lambda_j, \lambda_{j+1}, \dots, \lambda_m)$ and

$$\langle \lambda_{(j)} | = \langle \lambda_1 | \otimes \dots \otimes \langle -\lambda_j | \otimes \dots \otimes \langle \lambda_m |, \\ \langle \{\lambda, n\}_{(j)} | = \langle \lambda_{(j)} | \otimes \langle n | \exp \left\{ \frac{\Lambda(\lambda_{(j)})x}{2} (a^\dagger - a) \right\},$$

and

$$\Lambda(\lambda_{(j)}) = \lambda_1 + \dots + \lambda_{j-1} - \lambda_j + \lambda_{j+1} + \dots + \lambda_m = \Lambda(\lambda) - 2\lambda_j$$

and

$$E_n(\lambda) - E_{n'}(\lambda_{(j)}) = \omega \left\{ -\frac{x^2}{4} \Lambda(\lambda)^2 + n \right\} - \omega \left\{ -\frac{x^2}{4} \Lambda(\lambda_{(j)})^2 + n' \right\} \\ = \omega(n - n') + \omega x^2(1 - \lambda_j \Lambda(\lambda)).$$

In the following we set for simplicity $\phi_j = 0$ and

$$\Theta_j(t) \equiv g_2 \frac{\sin(\omega_j t)}{\omega_j} \quad \text{for } 1 \leq j \leq m, \quad (16)$$

then

$$H_F = \sum_{\lambda} \sum_{n, n'} \sum_{j=1}^m e^{i\left\{t\omega(n-n') + t\omega x^2(1-\lambda_j\Lambda(\lambda)) + 2\lambda_j\Theta_j(t)\right\}} \langle n | e^{\lambda_j x(a^\dagger - a)} | n' \rangle |\{\lambda, n\}\rangle \langle \{\lambda, n'\}_{(j)}|. \quad (17)$$

Here we divide H_F into two parts $H_F = H_F' + H_F''$ where

$$H_F' = \sum_{\lambda} \sum_n \sum_{j=1}^m e^{i\{t\omega x^2(1-\lambda_j\Lambda(\lambda))+2\lambda_j\Theta_j(t)\}} \langle n | e^{\lambda_j x(a^\dagger - a)} | n \rangle |\{\lambda, n\}\rangle \langle \{\lambda, n\}_{(j)}|, \quad (18)$$

and

$$H_F'' = \sum_{\lambda} \sum_{\substack{n, n' \\ n \neq n'}} \sum_{j=1}^m e^{i\{t\omega(n-n') + t\omega x^2(1-\lambda_j\Lambda(\lambda))+2\lambda_j\Theta_j(t)\}} \langle n | e^{\lambda_j x(a^\dagger - a)} | n' \rangle |\{\lambda, n\}\rangle \langle \{\lambda, n'\}_{(j)}|. \quad (19)$$

Noting $\langle n | e^{x(a^\dagger - a)} | n \rangle = \langle n | e^{-x(a^\dagger - a)} | n \rangle$ ($\lambda_j = \pm 1$) by the results in section 3 of [13],

H_F' can be written as

$$H_F' = \sum_n \langle n | e^{x(a^\dagger - a)} | n \rangle \left\{ \sum_{\lambda} \sum_{j=1}^m e^{i\{t\omega x^2(1-\lambda_j\Lambda(\lambda))+2\lambda_j\Theta_j(t)\}} |\{\lambda, n\}\rangle \langle \{\lambda, n\}_{(j)}| \right\}. \quad (20)$$

For us it is not easy to expand the equation above, so we consider the special case.

Case of $m=2$:

$$\begin{aligned} & \left\{ \sum_{\lambda} \sum_{j=1}^2 \dots \right\} \\ = & e^{i\{-t\omega x^2 + 2\Theta_1(t)\}} |\{1, 1, n\}\rangle \langle \{1, 1, n\}_{(1)}| + e^{i\{-t\omega x^2 + 2\Theta_2(t)\}} |\{1, 1, n\}\rangle \langle \{1, 1, n\}_{(2)}| + \\ & e^{i\{t\omega x^2 + 2\Theta_1(t)\}} |\{1, -1, n\}\rangle \langle \{1, -1, n\}_{(1)}| + e^{i\{t\omega x^2 - 2\Theta_2(t)\}} |\{1, -1, n\}\rangle \langle \{1, -1, n\}_{(2)}| + \\ & e^{i\{t\omega x^2 - 2\Theta_1(t)\}} |\{-1, 1, n\}\rangle \langle \{-1, 1, n\}_{(1)}| + e^{i\{t\omega x^2 + 2\Theta_2(t)\}} |\{-1, 1, n\}\rangle \langle \{-1, 1, n\}_{(2)}| + \\ & e^{i\{-t\omega x^2 - 2\Theta_1(t)\}} |\{-1, -1, n\}\rangle \langle \{-1, -1, n\}_{(1)}| + e^{i\{-t\omega x^2 - 2\Theta_2(t)\}} |\{-1, -1, n\}\rangle \langle \{-1, -1, n\}_{(2)}| \\ = & e^{i\{-t\omega x^2 + 2\Theta_1(t)\}} |\{1, 1, n\}\rangle \langle \{1, 1, n\}_{(1)}| + e^{i\{-t\omega x^2 + 2\Theta_2(t)\}} |\{1, 1, n\}\rangle \langle \{1, 1, n\}_{(2)}| + \\ & e^{i\{-t\omega x^2 - 2\Theta_2(t)\}} |\{-1, -1, n\}\rangle \langle \{-1, -1, n\}_{(2)}| + e^{i\{-t\omega x^2 - 2\Theta_1(t)\}} |\{-1, -1, n\}\rangle \langle \{-1, -1, n\}_{(1)}| + \\ & e^{i\{t\omega x^2 - 2\Theta_2(t)\}} |\{1, -1, n\}\rangle \langle \{1, -1, n\}_{(2)}| + e^{i\{t\omega x^2 + 2\Theta_1(t)\}} |\{1, -1, n\}\rangle \langle \{1, -1, n\}_{(1)}| + \\ & e^{i\{t\omega x^2 - 2\Theta_1(t)\}} |\{-1, 1, n\}\rangle \langle \{-1, 1, n\}_{(1)}| + e^{i\{t\omega x^2 + 2\Theta_2(t)\}} |\{-1, 1, n\}\rangle \langle \{-1, 1, n\}_{(2)}| \\ = & e^{i\{-t\omega x^2 + 2\Theta_1(t)\}} |\{1, 1, n\}\rangle \langle \{-1, 1, n\}| + e^{i\{-t\omega x^2 + 2\Theta_2(t)\}} |\{1, 1, n\}\rangle \langle \{1, -1, n\}| + \\ & e^{i\{-t\omega x^2 - 2\Theta_2(t)\}} |\{-1, -1, n\}\rangle \langle \{-1, 1, n\}| + e^{i\{-t\omega x^2 - 2\Theta_1(t)\}} |\{-1, -1, n\}\rangle \langle \{1, -1, n\}| + \\ & e^{i\{t\omega x^2 - 2\Theta_1(t)\}} |\{-1, 1, n\}\rangle \langle \{1, 1, n\}| + e^{i\{t\omega x^2 - 2\Theta_2(t)\}} |\{1, -1, n\}\rangle \langle \{1, 1, n\}| + \\ & e^{i\{t\omega x^2 + 2\Theta_2(t)\}} |\{-1, 1, n\}\rangle \langle \{-1, -1, n\}| + e^{i\{t\omega x^2 + 2\Theta_1(t)\}} |\{1, -1, n\}\rangle \langle \{-1, -1, n\}|, \end{aligned} \quad (21)$$

where let us once more remind

$$\begin{aligned}
|\{1, 1, n\}\rangle &= |1\rangle \otimes |1\rangle \otimes e^{-x(a^\dagger - a)}|n\rangle, \\
|\{1, -1, n\}\rangle &= |1\rangle \otimes |-1\rangle \otimes |n\rangle, \\
|\{-1, 1, n\}\rangle &= |-1\rangle \otimes |1\rangle \otimes |n\rangle, \\
|\{-1, -1, n\}\rangle &= |-1\rangle \otimes |-1\rangle \otimes e^{x(a^\dagger - a)}|n\rangle.
\end{aligned} \tag{22}$$

Here we want to solve the equation $i(d/dt)\Psi_0 = H_F'\Psi_0$ completely, however it is very hard (see for example [7], [10], [21], [22]). Therefore let us appeal to a perturbation theory. Now we use the well-known formula

$$e^{2i\lambda_j\Theta_j(t)} = \sum_{\alpha_j \in \mathbf{Z}} J_{\alpha_j}(2\lambda_j g_2/\omega_j) e^{i\alpha_j \omega_j t} = J_0(2g_2/\omega_j) + \sum_{\alpha_j \neq 0} J_{\alpha_j}(2\lambda_j g_2/\omega_j) e^{i\alpha_j \omega_j t}, \tag{23}$$

where $J_\alpha(x)$ are the Bessel functions. For a further simplicity we set $2g_2/\omega_j = \Gamma_j$.

Here we define a kind of extended Bell states from (22) ($|1\rangle \longleftrightarrow |0\rangle$, $|-1\rangle \longleftrightarrow |1\rangle$ in a usual manner)

$$\begin{aligned}
|\{\Phi_1, n\}\rangle &= \frac{1}{\sqrt{2}} (|\{1, 1, n\}\rangle + |\{-1, -1, n\}\rangle), \\
|\{\Phi_2, n\}\rangle &= \frac{1}{\sqrt{2}} (|\{1, 1, n\}\rangle - |\{-1, -1, n\}\rangle), \\
|\{\Phi_3, n\}\rangle &= \frac{1}{\sqrt{2}} (|\{-1, 1, n\}\rangle + |\{1, -1, n\}\rangle), \\
|\{\Phi_4, n\}\rangle &= \frac{1}{\sqrt{2}} (|\{-1, 1, n\}\rangle - |\{1, -1, n\}\rangle).
\end{aligned} \tag{24}$$

A comment is in order. We want to call these states (bases) Bell–Schrödinger cat states. Conversely we have

$$\begin{aligned}
|\{1, 1, n\}\rangle &= \frac{1}{\sqrt{2}} (|\{\Phi_1, n\}\rangle + |\{\Phi_2, n\}\rangle), \\
|\{-1, -1, n\}\rangle &= \frac{1}{\sqrt{2}} (|\{\Phi_1, n\}\rangle - |\{\Phi_2, n\}\rangle), \\
|\{-1, 1, n\}\rangle &= \frac{1}{\sqrt{2}} (|\{\Phi_3, n\}\rangle + |\{\Phi_4, n\}\rangle), \\
|\{1, -1, n\}\rangle &= \frac{1}{\sqrt{2}} (|\{\Phi_3, n\}\rangle - |\{\Phi_4, n\}\rangle).
\end{aligned} \tag{25}$$

From (14)

$$i\frac{d}{dt}\Psi_0 = \frac{\Delta}{2}H_F\Psi_0 = \frac{\Delta}{2}(H_F' + H_F'')\Psi_0.$$

However we are not interested in a transition from one state to an another one from the lesson obtaining one qubit case in [16], so we can remove H_F'' ($n \neq n'$) from the Hamiltonian. Next let us transform H_F' . From (21) and (23) if we set

$$\left\{ \sum_{\lambda} \sum_{j=1}^2 \cdots \right\} \equiv K_{0F}' + K_{1F}' :$$

then we obtain after some algebras

$$\begin{aligned} K_{0F}' = & (J_0(\Gamma_1) + J_0(\Gamma_2)) \left\{ e^{-it\omega x^2} |\{\Phi_1, n\}\rangle \langle \{\Phi_3, n\}| + e^{it\omega x^2} |\{\Phi_3, n\}\rangle \langle \{\Phi_1, n\}| \right\} + \\ & (J_0(\Gamma_1) - J_0(\Gamma_2)) \left\{ e^{-it\omega x^2} |\{\Phi_2, n\}\rangle \langle \{\Phi_4, n\}| + e^{it\omega x^2} |\{\Phi_4, n\}\rangle \langle \{\Phi_2, n\}| \right\}, \quad (26) \end{aligned}$$

and

$$\begin{aligned} K_{1F}' = & e^{-it\omega x^2} \sum_{\alpha \neq 0} \left\{ \frac{J_{\alpha}(\Gamma_1) + J_{\alpha}(-\Gamma_1)}{2} e^{it\alpha\omega_1} + \frac{J_{\alpha}(\Gamma_2) + J_{\alpha}(-\Gamma_2)}{2} e^{it\alpha\omega_2} \right\} |\{\Phi_1, n\}\rangle \langle \{\Phi_3, n\}| + \\ & e^{-it\omega x^2} \sum_{\alpha \neq 0} \left\{ \frac{J_{\alpha}(\Gamma_1) - J_{\alpha}(-\Gamma_1)}{2} e^{it\alpha\omega_1} - \frac{J_{\alpha}(\Gamma_2) - J_{\alpha}(-\Gamma_2)}{2} e^{it\alpha\omega_2} \right\} |\{\Phi_1, n\}\rangle \langle \{\Phi_4, n\}| + \\ & e^{-it\omega x^2} \sum_{\alpha \neq 0} \left\{ \frac{J_{\alpha}(\Gamma_1) - J_{\alpha}(-\Gamma_1)}{2} e^{it\alpha\omega_1} + \frac{J_{\alpha}(\Gamma_2) - J_{\alpha}(-\Gamma_2)}{2} e^{it\alpha\omega_2} \right\} |\{\Phi_2, n\}\rangle \langle \{\Phi_3, n\}| + \\ & e^{-it\omega x^2} \sum_{\alpha \neq 0} \left\{ \frac{J_{\alpha}(\Gamma_1) + J_{\alpha}(-\Gamma_1)}{2} e^{it\alpha\omega_1} - \frac{J_{\alpha}(\Gamma_2) + J_{\alpha}(-\Gamma_2)}{2} e^{it\alpha\omega_2} \right\} |\{\Phi_2, n\}\rangle \langle \{\Phi_4, n\}| + \\ & e^{it\omega x^2} \sum_{\alpha \neq 0} \left\{ \frac{J_{\alpha}(-\Gamma_1) + J_{\alpha}(\Gamma_1)}{2} e^{it\alpha\omega_1} + \frac{J_{\alpha}(-\Gamma_2) + J_{\alpha}(\Gamma_2)}{2} e^{it\alpha\omega_2} \right\} |\{\Phi_3, n\}\rangle \langle \{\Phi_1, n\}| + \\ & e^{it\omega x^2} \sum_{\alpha \neq 0} \left\{ \frac{J_{\alpha}(-\Gamma_1) - J_{\alpha}(\Gamma_1)}{2} e^{it\alpha\omega_1} - \frac{J_{\alpha}(-\Gamma_2) - J_{\alpha}(\Gamma_2)}{2} e^{it\alpha\omega_2} \right\} |\{\Phi_4, n\}\rangle \langle \{\Phi_1, n\}| + \\ & e^{it\omega x^2} \sum_{\alpha \neq 0} \left\{ \frac{J_{\alpha}(-\Gamma_1) - J_{\alpha}(\Gamma_1)}{2} e^{it\alpha\omega_1} + \frac{J_{\alpha}(-\Gamma_2) - J_{\alpha}(\Gamma_2)}{2} e^{it\alpha\omega_2} \right\} |\{\Phi_3, n\}\rangle \langle \{\Phi_2, n\}| + \\ & e^{it\omega x^2} \sum_{\alpha \neq 0} \left\{ \frac{J_{\alpha}(-\Gamma_1) + J_{\alpha}(\Gamma_1)}{2} e^{it\alpha\omega_1} - \frac{J_{\alpha}(-\Gamma_2) + J_{\alpha}(\Gamma_2)}{2} e^{it\alpha\omega_2} \right\} |\{\Phi_4, n\}\rangle \langle \{\Phi_2, n\}|, \quad (27) \end{aligned}$$

where we have used the basic relations

$$J_{\alpha}(-x) = (-1)^{\alpha} J_{\alpha}(x) = J_{-\alpha}(x).$$

Therefore

$$i\frac{d}{dt}\Psi_0 = \frac{\Delta}{2}H_F'\Psi_0 = \sum_n \left\{ \frac{\Delta}{2}\langle n|e^{x(a^\dagger-a)}|n\rangle K_{0F}' + \frac{\Delta}{2}\langle n|e^{x(a^\dagger-a)}|n\rangle K_{1F}' \right\} \Psi_0. \quad (28)$$

In the following we treat $\frac{\Delta}{2}\langle n|e^{x(a^\dagger-a)}|n\rangle K_{0F}'$ a unperturbed Hamiltonian and the remaining a perturbed one. If we set for simplicity

$$\begin{aligned} E_{\Delta,n,+} &= \frac{\Delta}{2}\langle n|e^{x(a^\dagger-a)}|n\rangle (J_0(\Gamma_1) + J_0(\Gamma_2)), \\ E_{\Delta,n,-} &= \frac{\Delta}{2}\langle n|e^{x(a^\dagger-a)}|n\rangle (J_0(\Gamma_1) - J_0(\Gamma_2)), \end{aligned} \quad (29)$$

then it is easy to solve

$$\begin{aligned} i\frac{d}{dt}\Psi_0 &= \sum_n \frac{\Delta}{2} \left\{ \langle n|e^{x(a^\dagger-a)}|n\rangle K_{0F}' \right\} \Psi_0 \\ &= \sum_n \left[E_{\Delta,n,+} \left\{ e^{-it\omega x^2} |\{\Phi_1, n\}\rangle \langle \{\Phi_3, n\}| + e^{it\omega x^2} |\{\Phi_3, n\}\rangle \langle \{\Phi_1, n\}| \right\} + \right. \\ &\quad \left. E_{\Delta,n,-} \left\{ e^{-it\omega x^2} |\{\Phi_2, n\}\rangle \langle \{\Phi_4, n\}| + e^{it\omega x^2} |\{\Phi_4, n\}\rangle \langle \{\Phi_2, n\}| \right\} \right] \Psi_0, \end{aligned}$$

see Appendix.

As an application to Quantum Computation it is sufficient for us to consider one excited state. By making use of the method of constant variation again we can set $\Psi_0(t)$ as

$$\begin{aligned} \Psi_0(t) &= (u_{n,11}c_{n,1}(t) + u_{n,13}c_{n,3}(t))|\{\Phi_1, n\}\rangle + (u_{n,22}c_{n,2}(t) + u_{n,24}c_{n,4}(t))|\{\Phi_2, n\}\rangle + \\ &\quad (u_{n,31}c_{n,1}(t) + u_{n,33}c_{n,3}(t))|\{\Phi_3, n\}\rangle + (u_{n,42}c_{n,2}(t) + u_{n,44}c_{n,4}(t))|\{\Phi_4, n\}\rangle \end{aligned} \quad (30)$$

for some fixed n , where

$$U_n(t) = \begin{pmatrix} u_{n,11} & u_{n,13} \\ u_{n,31} & u_{n,33} \end{pmatrix} = \begin{pmatrix} 1 & \\ & e^{it\omega x^2} \end{pmatrix} \exp \left\{ -it \begin{pmatrix} 0 & E_{\Delta,n,+} \\ E_{\Delta,n,+} & \omega x^2 \end{pmatrix} \right\}, \quad (31)$$

$$V_n(t) = \begin{pmatrix} u_{n,22} & u_{n,24} \\ u_{n,42} & u_{n,44} \end{pmatrix} = \begin{pmatrix} 1 & \\ & e^{it\omega x^2} \end{pmatrix} \exp \left\{ -it \begin{pmatrix} 0 & E_{\Delta,n,-} \\ E_{\Delta,n,-} & \omega x^2 \end{pmatrix} \right\} \quad (32)$$

from the appendix.

Then substituting (30) into (28) and some tedious calculations lead us to

$$i \frac{d}{dt} \begin{pmatrix} c_{n,1} \\ c_{n,3} \\ c_{n,2} \\ c_{n,4} \end{pmatrix} = \begin{pmatrix} u_{n,11} & u_{n,13} & & \\ u_{n,31} & u_{n,33} & & \\ & & u_{n,22} & u_{n,24} \\ & & u_{n,42} & u_{n,44} \end{pmatrix}^{-1} \begin{pmatrix} 0 & A & 0 & B \\ \bar{A} & 0 & \bar{C} & 0 \\ 0 & C & 0 & D \\ \bar{B} & 0 & \bar{D} & 0 \end{pmatrix} \times \begin{pmatrix} u_{n,11} & u_{n,13} & & \\ u_{n,31} & u_{n,33} & & \\ & & u_{n,22} & u_{n,24} \\ & & u_{n,42} & u_{n,44} \end{pmatrix} \begin{pmatrix} c_{n,1} \\ c_{n,3} \\ c_{n,2} \\ c_{n,4} \end{pmatrix} \quad (33)$$

with

$$\begin{aligned} A &= e^{-it\omega x^2} \frac{\Delta}{2} \langle n | e^{x(a^\dagger - a)} | n \rangle \sum_{\alpha \neq 0} \left\{ \frac{J_\alpha(\Gamma_1) + J_\alpha(-\Gamma_1)}{2} e^{it\alpha\omega_1} + \frac{J_\alpha(\Gamma_2) + J_\alpha(-\Gamma_2)}{2} e^{it\alpha\omega_2} \right\} \\ &\equiv e^{-it\omega x^2} A_0, \\ B &= e^{-it\omega x^2} \frac{\Delta}{2} \langle n | e^{x(a^\dagger - a)} | n \rangle \sum_{\alpha \neq 0} \left\{ \frac{J_\alpha(\Gamma_1) - J_\alpha(-\Gamma_1)}{2} e^{it\alpha\omega_1} - \frac{J_\alpha(\Gamma_2) - J_\alpha(-\Gamma_2)}{2} e^{it\alpha\omega_2} \right\} \\ &\equiv e^{-it\omega x^2} B_0, \\ C &= e^{-it\omega x^2} \frac{\Delta}{2} \langle n | e^{x(a^\dagger - a)} | n \rangle \sum_{\alpha \neq 0} \left\{ \frac{J_\alpha(\Gamma_1) - J_\alpha(-\Gamma_1)}{2} e^{it\alpha\omega_1} + \frac{J_\alpha(\Gamma_2) - J_\alpha(-\Gamma_2)}{2} e^{it\alpha\omega_2} \right\} \\ &\equiv e^{-it\omega x^2} C_0, \\ D &= e^{-it\omega x^2} \frac{\Delta}{2} \langle n | e^{x(a^\dagger - a)} | n \rangle \sum_{\alpha \neq 0} \left\{ \frac{J_\alpha(\Gamma_1) + J_\alpha(-\Gamma_1)}{2} e^{it\alpha\omega_1} - \frac{J_\alpha(\Gamma_2) + J_\alpha(-\Gamma_2)}{2} e^{it\alpha\omega_2} \right\} \\ &\equiv e^{-it\omega x^2} D_0. \end{aligned} \quad (34)$$

We note that $\bar{A}_0 = A_0$, $\bar{D}_0 = D_0$ and $\bar{B}_0 = -B_0$, $\bar{C}_0 = -C_0$ (due to the fact $J_{-\alpha}(x) = J_\alpha(-x)$).

The above matrix equation decomposes into two parts

$$i \frac{d}{dt} \begin{pmatrix} c_{n,1} \\ c_{n,3} \end{pmatrix} = \begin{pmatrix} u_{n,11} & u_{n,13} \\ u_{n,31} & u_{n,33} \end{pmatrix}^{-1} \begin{pmatrix} 0 & A \\ \bar{A} & 0 \end{pmatrix} \begin{pmatrix} u_{n,11} & u_{n,13} \\ u_{n,31} & u_{n,33} \end{pmatrix} \begin{pmatrix} c_{n,1} \\ c_{n,3} \end{pmatrix} + \begin{pmatrix} u_{n,11} & u_{n,13} \\ u_{n,31} & u_{n,33} \end{pmatrix}^{-1} \begin{pmatrix} 0 & B \\ \bar{C} & 0 \end{pmatrix} \begin{pmatrix} u_{n,22} & u_{n,24} \\ u_{n,42} & u_{n,44} \end{pmatrix} \begin{pmatrix} c_{n,2} \\ c_{n,4} \end{pmatrix},$$

$$i \frac{d}{dt} \begin{pmatrix} c_{n,2} \\ c_{n,4} \end{pmatrix} = \begin{pmatrix} u_{n,22} & u_{n,24} \\ u_{n,42} & u_{n,44} \end{pmatrix}^{-1} \begin{pmatrix} 0 & C \\ \bar{B} & 0 \end{pmatrix} \begin{pmatrix} u_{n,11} & u_{n,13} \\ u_{n,31} & u_{n,33} \end{pmatrix} \begin{pmatrix} c_{n,1} \\ c_{n,3} \end{pmatrix} + \\ \begin{pmatrix} u_{n,22} & u_{n,24} \\ u_{n,42} & u_{n,44} \end{pmatrix}^{-1} \begin{pmatrix} 0 & D \\ \bar{D} & 0 \end{pmatrix} \begin{pmatrix} u_{n,22} & u_{n,24} \\ u_{n,42} & u_{n,44} \end{pmatrix} \begin{pmatrix} c_{n,2} \\ c_{n,4} \end{pmatrix}.$$

By the way, from (31), (32) and (34)

$$i \frac{d}{dt} \begin{pmatrix} c_{n,1} \\ c_{n,3} \end{pmatrix} = \\ A_0 \exp \left\{ it \begin{pmatrix} 0 & E_{\Delta,n,+} \\ E_{\Delta,n,+} & \omega x^2 \end{pmatrix} \right\} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \exp \left\{ -it \begin{pmatrix} 0 & E_{\Delta,n,+} \\ E_{\Delta,n,+} & \omega x^2 \end{pmatrix} \right\} \begin{pmatrix} c_{n,1} \\ c_{n,3} \end{pmatrix} + \\ \exp \left\{ it \begin{pmatrix} 0 & E_{\Delta,n,+} \\ E_{\Delta,n,+} & \omega x^2 \end{pmatrix} \right\} \begin{pmatrix} 0 & B_0 \\ \bar{C}_0 & 0 \end{pmatrix} \exp \left\{ -it \begin{pmatrix} 0 & E_{\Delta,n,-} \\ E_{\Delta,n,-} & \omega x^2 \end{pmatrix} \right\} \begin{pmatrix} c_{n,2} \\ c_{n,4} \end{pmatrix}, \quad (35)$$

$$i \frac{d}{dt} \begin{pmatrix} c_{n,2} \\ c_{n,4} \end{pmatrix} = \\ \exp \left\{ it \begin{pmatrix} 0 & E_{\Delta,n,-} \\ E_{\Delta,n,-} & \omega x^2 \end{pmatrix} \right\} \begin{pmatrix} 0 & C_0 \\ \bar{B}_0 & 0 \end{pmatrix} \exp \left\{ -it \begin{pmatrix} 0 & E_{\Delta,n,+} \\ E_{\Delta,n,+} & \omega x^2 \end{pmatrix} \right\} \begin{pmatrix} c_{n,1} \\ c_{n,3} \end{pmatrix} + \\ D_0 \exp \left\{ it \begin{pmatrix} 0 & E_{\Delta,n,-} \\ E_{\Delta,n,-} & \omega x^2 \end{pmatrix} \right\} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \exp \left\{ -it \begin{pmatrix} 0 & E_{\Delta,n,-} \\ E_{\Delta,n,-} & \omega x^2 \end{pmatrix} \right\} \begin{pmatrix} c_{n,2} \\ c_{n,4} \end{pmatrix}. \quad (36)$$

At this stage we can set several resonance conditions and obtain corresponding equations and solutions by making use of the rotating wave approximation like in [15], [16].

For example, let us consider the term

$$\exp \left\{ it \begin{pmatrix} 0 & E_{\Delta,n,+} \\ E_{\Delta,n,+} & \omega x^2 \end{pmatrix} \right\} \begin{pmatrix} 0 & B_0 \\ \bar{C}_0 & 0 \end{pmatrix} \exp \left\{ -it \begin{pmatrix} 0 & E_{\Delta,n,-} \\ E_{\Delta,n,-} & \omega x^2 \end{pmatrix} \right\},$$

then from (57) in Appendix

$$P_{n,+} \begin{pmatrix} e^{it\mu_{n,+}} & \\ & e^{it\nu_{n,+}} \end{pmatrix} P_{n,+}^{-1} \begin{pmatrix} 0 & B_0 \\ \bar{C}_0 & 0 \end{pmatrix} P_{n,-} \begin{pmatrix} e^{-it\mu_{n,-}} & \\ & e^{-it\nu_{n,-}} \end{pmatrix} P_{n,-}^{-1}.$$

Here for example,

$$e^{it(\mu_{n,+}-\mu_{n,-})} P_{n,+} \begin{pmatrix} 1 & \\ & e^{it(\nu_{n,+}-\mu_{n,+})} \end{pmatrix} P_{n,+}^{-1} \begin{pmatrix} 0 & B_0 \\ \bar{C}_0 & 0 \end{pmatrix} P_{n,-} \begin{pmatrix} 1 & \\ & e^{-it(\nu_{n,-}-\mu_{n,-})} \end{pmatrix} P_{n,-}^{-1}$$

for

$$B_0 = \frac{\Delta}{2} \langle n | e^{x(a^\dagger - a)} | n \rangle \sum_{\alpha \neq 0} \left\{ \frac{J_\alpha(\Gamma_1) - J_\alpha(-\Gamma_1)}{2} e^{it\alpha\omega_1} - \frac{J_\alpha(\Gamma_2) - J_\alpha(-\Gamma_2)}{2} e^{it\alpha\omega_2} \right\},$$

$$\bar{C}_0 = \frac{\Delta}{2} \langle n | e^{x(a^\dagger - a)} | n \rangle \sum_{\alpha \neq 0} \left\{ \frac{J_\alpha(-\Gamma_1) - J_\alpha(\Gamma_1)}{2} e^{it\alpha\omega_1} - \frac{J_\alpha(\Gamma_2) - J_\alpha(-\Gamma_2)}{2} e^{it\alpha\omega_2} \right\}.$$

Now we set the resonance condition

$$\alpha\omega_2 + \mu_{n,+} - \mu_{n,-} = 0 \iff (-\alpha)\omega_2 + \mu_{n,-} - \mu_{n,+} = 0 \quad (37)$$

for some $\alpha \neq 0$, then we can neglect the remaining terms by the rotating wave approximation. That is, we obtain the time independent matrix

$$-\frac{\Delta}{2} \langle n | e^{x(a^\dagger - a)} | n \rangle \frac{J_\alpha(\Gamma_2) - J_\alpha(-\Gamma_2)}{2} P_{n,+} \begin{pmatrix} 1 & \\ & 0 \end{pmatrix} P_{n,+}^{-1} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} P_{n,-} \begin{pmatrix} 1 & \\ & 0 \end{pmatrix} P_{n,-}^{-1},$$

so we have

$$i \frac{d}{dt} \begin{pmatrix} c_{n,1} \\ c_{n,3} \end{pmatrix} = -\frac{\Delta}{2} \langle n | e^{x(a^\dagger - a)} | n \rangle \frac{J_\alpha(\Gamma_2) - J_\alpha(-\Gamma_2)}{2} \times$$

$$P_{n,+} \begin{pmatrix} 1 & \\ & 0 \end{pmatrix} P_{n,+}^{-1} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} P_{n,-} \begin{pmatrix} 1 & \\ & 0 \end{pmatrix} P_{n,-}^{-1} \begin{pmatrix} c_{n,2} \\ c_{n,4} \end{pmatrix},$$

$$i \frac{d}{dt} \begin{pmatrix} c_{n,2} \\ c_{n,4} \end{pmatrix} = -\frac{\Delta}{2} \langle n | e^{x(a^\dagger - a)} | n \rangle \frac{J_\alpha(\Gamma_2) - J_\alpha(-\Gamma_2)}{2} \times$$

$$P_{n,-} \begin{pmatrix} 1 & \\ & 0 \end{pmatrix} P_{n,-}^{-1} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} P_{n,+} \begin{pmatrix} 1 & \\ & 0 \end{pmatrix} P_{n,+}^{-1} \begin{pmatrix} c_{n,1} \\ c_{n,3} \end{pmatrix},$$

where we have used the relation

$$\frac{J_{-\alpha}(\Gamma_2) - J_{-\alpha}(-\Gamma_2)}{2} = -\frac{J_\alpha(\Gamma_2) - J_\alpha(-\Gamma_2)}{2}.$$

After some algebras we obtain

$$\begin{aligned}
i \frac{d}{dt} \begin{pmatrix} c_{n,1} \\ c_{n,3} \end{pmatrix} &= -\frac{\Delta}{2} \langle n | e^{x(a^\dagger - a)} | n \rangle \frac{J_\alpha(\Gamma_2) - J_\alpha(-\Gamma_2)}{2} \times \\
&\quad \frac{E_{\Delta,n,+}\mu_{n,-} + E_{\Delta,n,-}\mu_{n,+}}{(E_{\Delta,n,+}^2 + \mu_{n,+}^2)(E_{\Delta,n,-}^2 + \mu_{n,-}^2)} \begin{pmatrix} E_{\Delta,n,+}E_{\Delta,n,-} & E_{\Delta,n,+}\mu_{n,-} \\ \mu_{n,+}E_{\Delta,n,-} & \mu_{n,+}\mu_{n,-} \end{pmatrix} \begin{pmatrix} c_{n,2} \\ c_{n,4} \end{pmatrix} \quad (38) \\
i \frac{d}{dt} \begin{pmatrix} c_{n,2} \\ c_{n,4} \end{pmatrix} &= -\frac{\Delta}{2} \langle n | e^{x(a^\dagger - a)} | n \rangle \frac{J_\alpha(\Gamma_2) - J_\alpha(-\Gamma_2)}{2} \times \\
&\quad \frac{E_{\Delta,n,+}\mu_{n,-} + E_{\Delta,n,-}\mu_{n,+}}{(E_{\Delta,n,+}^2 + \mu_{n,+}^2)(E_{\Delta,n,-}^2 + \mu_{n,-}^2)} \begin{pmatrix} E_{\Delta,n,+}E_{\Delta,n,-} & \mu_{n,+}E_{\Delta,n,-} \\ E_{\Delta,n,+}\mu_{n,-} & \mu_{n,+}\mu_{n,-} \end{pmatrix} \begin{pmatrix} c_{n,1} \\ c_{n,3} \end{pmatrix} \quad (39)
\end{aligned}$$

from the appendix.

If we prepare the vector notations

$$\mathbf{c}_{od} = \begin{pmatrix} c_{n,1} \\ c_{n,3} \end{pmatrix}, \quad \mathbf{c}_{ev} = \begin{pmatrix} c_{n,2} \\ c_{n,4} \end{pmatrix},$$

then the equation above can be written as

$$i \frac{d}{dt} \begin{pmatrix} \mathbf{c}_{od} \\ \mathbf{c}_{ev} \end{pmatrix} = -\frac{\mathcal{R}}{2} \begin{pmatrix} & K \\ K^T & \end{pmatrix} \begin{pmatrix} \mathbf{c}_{od} \\ \mathbf{c}_{ev} \end{pmatrix} \quad (40)$$

with

$$\mathcal{R} = \frac{\Delta}{2} \langle n | e^{x(a^\dagger - a)} | n \rangle \frac{J_\alpha(\Gamma_2) - J_\alpha(-\Gamma_2)}{2} \frac{E_{\Delta,n,+}\mu_{n,-} + E_{\Delta,n,-}\mu_{n,+}}{\sqrt{(E_{\Delta,n,+}^2 + \mu_{n,+}^2)(E_{\Delta,n,-}^2 + \mu_{n,-}^2)}}$$

and

$$K = \frac{1}{\sqrt{(E_{\Delta,n,+}^2 + \mu_{n,+}^2)(E_{\Delta,n,-}^2 + \mu_{n,-}^2)}} \begin{pmatrix} E_{\Delta,n,+}E_{\Delta,n,-} & E_{\Delta,n,+}\mu_{n,-} \\ \mu_{n,+}E_{\Delta,n,-} & \mu_{n,+}\mu_{n,-} \end{pmatrix}.$$

The (formal) solution is easily obtained to become

$$\begin{pmatrix} \mathbf{c}_{od}(t) \\ \mathbf{c}_{ev}(t) \end{pmatrix} = \exp \left\{ \frac{i\mathcal{R}t}{2} \begin{pmatrix} & K \\ K^T & \end{pmatrix} \right\} \begin{pmatrix} \mathbf{c}_{od}(0) \\ \mathbf{c}_{ev}(0) \end{pmatrix}. \quad (41)$$

Next let us calculate the unitary transformation of (41). It is easy to see

$$K = \frac{1}{\sqrt{(E_{\Delta,n,+}^2 + \mu_{n,+}^2)(E_{\Delta,n,-}^2 + \mu_{n,-}^2)}} \begin{pmatrix} E_{\Delta,n,+} \\ \mu_{n,+} \end{pmatrix} (E_{\Delta,n,-}, \mu_{n,-}), \quad (42)$$

$$K^T = \frac{1}{\sqrt{(E_{\Delta,n,+}^2 + \mu_{n,+}^2)(E_{\Delta,n,-}^2 + \mu_{n,-}^2)}} \begin{pmatrix} E_{\Delta,n,-} \\ \mu_{n,-} \end{pmatrix} (E_{\Delta,n,+}, \mu_{n,+}), \quad (43)$$

so

$$\begin{aligned} K^T K &= \frac{1}{E_{\Delta,n,-}^2 + \mu_{n,-}^2} \begin{pmatrix} E_{\Delta,n,-} \\ \mu_{n,-} \end{pmatrix} (E_{\Delta,n,-}, \mu_{n,-}) \\ &= \begin{pmatrix} E_{\Delta,n,-} \\ \mu_{n,-} \end{pmatrix} (E_{\Delta,n,-}^2 + \mu_{n,-}^2)^{-1} (E_{\Delta,n,-}, \mu_{n,-}), \\ K K^T &= \frac{1}{E_{\Delta,n,+}^2 + \mu_{n,+}^2} \begin{pmatrix} E_{\Delta,n,+} \\ \mu_{n,+} \end{pmatrix} (E_{\Delta,n,+}, \mu_{n,+}) \\ &= \begin{pmatrix} E_{\Delta,n,+} \\ \mu_{n,+} \end{pmatrix} (E_{\Delta,n,+}^2 + \mu_{n,+}^2)^{-1} (E_{\Delta,n,+}, \mu_{n,+}). \end{aligned}$$

Namely, $K^T K$ and $K K^T$ are just projection matrices and moreover satisfy

$$(K K^T)^n = K K^T, \quad (K^T K)^n = K^T K, \quad K K^T K = K, \quad K^T K K^T = K^T$$

for $n \geq 1$. Therefore after long algebras using the relations above we have

$$\begin{aligned} \exp \left\{ \frac{i\mathcal{R}t}{2} \begin{pmatrix} & K \\ K^T & \end{pmatrix} \right\} = \\ \begin{pmatrix} \mathbf{1} - K K^T + \cos\left(\frac{\mathcal{R}t}{2}\right) K K^T & i\sin\left(\frac{\mathcal{R}t}{2}\right) K \\ i\sin\left(\frac{\mathcal{R}t}{2}\right) K^T & \mathbf{1} - K^T K + \cos\left(\frac{\mathcal{R}t}{2}\right) K^T K \end{pmatrix}, \end{aligned} \quad (44)$$

so the solution is explicitly

$$\begin{pmatrix} \mathbf{c}_{od}(t) \\ \mathbf{c}_{ev}(t) \end{pmatrix} = \begin{pmatrix} \mathbf{1} - K K^T + \cos\left(\frac{\mathcal{R}t}{2}\right) K K^T & i\sin\left(\frac{\mathcal{R}t}{2}\right) K \\ i\sin\left(\frac{\mathcal{R}t}{2}\right) K^T & \mathbf{1} - K^T K + \cos\left(\frac{\mathcal{R}t}{2}\right) K^T K \end{pmatrix} \begin{pmatrix} \mathbf{c}_{od}(0) \\ \mathbf{c}_{ev}(0) \end{pmatrix} \quad (45)$$

with (42) and (43).

This is one of unitary transformations that we are looking for in Quantum Computation. We can also obtain another ones by setting different resonance conditions like (37) (we leave them to the readers).

Lastly let us summarize our result. By (4) the Hamiltonian becomes

$$H_L = \sum_{j=1}^m \left\{ \frac{\Delta}{2} \sigma_3^{(j)} + g_2 \cos(\omega_j t + \phi_j) \sigma_1^{(j)} \right\} \otimes \mathbf{1}_L \quad (46)$$

if there is no photon interaction, so the dynamics of each qubit space is independently determined by

$$H_{Lj} = \frac{\Delta}{2} \sigma_3^{(j)} + g_2 \cos(\omega_j t + \phi_j) \sigma_1^{(j)}. \quad (47)$$

Therefore the total space of m -qubits is just


$$\mathbf{C}^2 \otimes \dots \otimes \mathbf{C}^2 \otimes \dots \otimes \mathbf{C}^2 \quad \text{where} \quad \mathbf{C}^2 = \text{Vect}_{\mathbf{C}}\{|0\rangle, |1\rangle\},$$

and unitary transformations of each qubit space are obtained by manipulating laser fields. However to solve the Schrödinger equation (neglecting the suffix)

$$i \frac{d}{dt} \Psi = \left\{ \frac{\Delta}{2} \sigma_3 + g_2 \cos(\omega t + \phi) \sigma_1 \right\} \Psi$$

is not so easy, see for example [10].

When $m = 2$, the interaction (driving by the photon) between two qubits is given by unitary transformations like (45) $\dots$ controlled unitary gates including the controlled NOT.




This is just our scenario of Quantum Computation in the strong coupling regime.

Case of $m=3$:

We present a very important

Problem : Let us consider three atoms in a cavity. How can we construct C-NOT (or C-Unitary) operations for any two atoms among them ?

See the following pictures :



These constructions are very crucial in realizing quantum logic gates, however we have not seen such constructions in any references. We will attack this problem in a forthcoming paper.

In this paper we treated the two-atoms case in a cavity QED and constructed unitary transformations by making use of the rotating wave approximation under new resonance conditions containing the Bessel functions.

These will be applied to construct several quantum logic gates in Quantum Computation. Moreover we would like to treat a general case, which is at the present not easy due to some technical reasons.

By the way, according to increase of the number of atoms (we are expecting at least $m = 100$ in the realistic quantum computation) we meet a very severe problem called Decoherence, see for example [9] and its references. We don't know how to control this. One way protecting against this may be to deal with N-level system (then we can reduce the number of atoms in a cavity). A generalization of the model to N-level system (see for example [14], [19], [20], [24]) is now under consideration and will be published in a separate paper.

Acknowledgment. The author wishes to thank Marco Frasca for his important suggestions. He also wishes to thank the graduate students Kyoko Higashida, Ryosuke Kato and Yukako Wada for some help.

Appendix : Some Useful Formulas

In this appendix we solve the following equation

$$i\frac{d}{dt}\psi = \alpha H\psi, \quad (48)$$

where α is a constant and

$$H = e^{-i\theta t}|1\rangle\langle -1| + e^{i\theta t}|-1\rangle\langle 1| \quad \text{and} \quad \psi = a(t)|1\rangle + b(t)|-1\rangle. \quad (49)$$

Then it is easy to get a matrix equation on $\{a, b\}$

$$i\frac{d}{dt}\begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 & \alpha e^{-i\theta t} \\ \alpha e^{i\theta t} & 0 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} \Longleftrightarrow i\frac{d}{dt}\tilde{\psi} = \tilde{H}\tilde{\psi}. \quad (50)$$

Noting the decomposition

$$\begin{pmatrix} 0 & \alpha e^{-i\theta t} \\ \alpha e^{i\theta t} & 0 \end{pmatrix} = \begin{pmatrix} 1 & \\ & e^{i\theta t} \end{pmatrix} \begin{pmatrix} 0 & \alpha \\ \alpha & 0 \end{pmatrix} \begin{pmatrix} 1 & \\ & e^{-i\theta t} \end{pmatrix},$$

then the equation becomes

$$i \frac{d}{dt} \hat{\psi} = \begin{pmatrix} 0 & \alpha \\ \alpha & \theta \end{pmatrix} \hat{\psi} \quad \text{for} \quad \hat{\psi} \equiv \begin{pmatrix} 1 & \\ & e^{-i\theta t} \end{pmatrix} \tilde{\psi}.$$

The solution is easily obtained to become

$$\begin{pmatrix} a \\ b \end{pmatrix} = U(t) \begin{pmatrix} a_0 \\ b_0 \end{pmatrix} \quad (51)$$

where $(a_0, b_0)^T$ is a constant vector and

$$U(t) = \begin{pmatrix} 1 & \\ & e^{i\theta t} \end{pmatrix} \exp \left\{ -it \begin{pmatrix} 0 & \alpha \\ \alpha & \theta \end{pmatrix} \right\} \Rightarrow i \frac{d}{dt} U = \tilde{H} U. \quad (52)$$

If we set

$$U(t) = \begin{pmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{pmatrix} \quad (53)$$

(2 corresponds to -1) then ψ above can be written as

$$\psi = (u_{11}a_0 + u_{12}b_0)|1\rangle + (u_{21}a_0 + u_{22}b_0)|-1\rangle \quad (54)$$

with constants $\{a_0, b_0\}$.

In the method of constant variation in the text we change like $a_0 \longrightarrow a_0(t)$ and $b_0 \longrightarrow b_0(t)$.

Let us make some comments. For

$$A = \begin{pmatrix} 0 & \alpha \\ \alpha & \theta \end{pmatrix} \quad (55)$$

we can easily diagonalize A as follows :

$$A = \begin{pmatrix} \frac{\alpha}{\sqrt{\alpha^2 + \mu^2}} & \frac{\alpha}{\sqrt{\alpha^2 + \nu^2}} \\ \frac{\mu}{\sqrt{\alpha^2 + \mu^2}} & \frac{\nu}{\sqrt{\alpha^2 + \nu^2}} \end{pmatrix} \begin{pmatrix} \mu & \\ & \nu \end{pmatrix} \begin{pmatrix} \frac{\alpha}{\sqrt{\alpha^2 + \mu^2}} & \frac{\alpha}{\sqrt{\alpha^2 + \nu^2}} \\ \frac{\mu}{\sqrt{\alpha^2 + \mu^2}} & \frac{\nu}{\sqrt{\alpha^2 + \nu^2}} \end{pmatrix}^{-1} \quad (56)$$

where

$$\mu = \frac{1}{2}(\theta + \sqrt{\theta^2 + 4\alpha^2}), \quad \nu = \frac{1}{2}(\theta - \sqrt{\theta^2 + 4\alpha^2}).$$

Therefore we obtain

$$Q(t) \equiv e^{-itA} = \begin{pmatrix} \frac{\alpha}{\sqrt{\alpha^2 + \mu^2}} & \frac{\alpha}{\sqrt{\alpha^2 + \nu^2}} \\ \frac{\mu}{\sqrt{\alpha^2 + \mu^2}} & \frac{\nu}{\sqrt{\alpha^2 + \nu^2}} \end{pmatrix} \begin{pmatrix} e^{-it\mu} & \\ & e^{-it\nu} \end{pmatrix} \begin{pmatrix} \frac{\alpha}{\sqrt{\alpha^2 + \mu^2}} & \frac{\alpha}{\sqrt{\alpha^2 + \nu^2}} \\ \frac{\mu}{\sqrt{\alpha^2 + \mu^2}} & \frac{\nu}{\sqrt{\alpha^2 + \nu^2}} \end{pmatrix}^{-1} \quad (57)$$

For the simplicity we set

$$P = \begin{pmatrix} \frac{\alpha}{\sqrt{\alpha^2 + \mu^2}} & \frac{\alpha}{\sqrt{\alpha^2 + \nu^2}} \\ \frac{\mu}{\sqrt{\alpha^2 + \mu^2}} & \frac{\nu}{\sqrt{\alpha^2 + \nu^2}} \end{pmatrix} \in O(2). \quad (58)$$

In the text this is used as

$$P_{n,\sigma} = \begin{pmatrix} \frac{E_{\Delta,n,\sigma}}{\sqrt{E_{\Delta,n,\sigma}^2 + \mu_{n,\sigma}^2}} & \frac{E_{\Delta,n,\sigma}}{\sqrt{E_{\Delta,n,\sigma}^2 + \nu_{n,\sigma}^2}} \\ \frac{\mu_{n,\sigma}}{\sqrt{E_{\Delta,n,\sigma}^2 + \mu_{n,\sigma}^2}} & \frac{\nu_{n,\sigma}}{\sqrt{E_{\Delta,n,\sigma}^2 + \nu_{n,\sigma}^2}} \end{pmatrix} \quad \text{for } \sigma = \pm \quad (59)$$

with

$$\mu_{n,\sigma} = \frac{1}{2}(\gamma + \sqrt{\gamma^2 + 4E_{\Delta,n,\sigma}^2}), \quad \nu_{n,\sigma} = \frac{1}{2}(\gamma - \sqrt{\gamma^2 + 4E_{\Delta,n,\sigma}^2}) \quad (60)$$

for $\gamma = \omega x^2$. Then

$$P_{n,+}^T \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} P_{n,-} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}, \quad (61)$$

where

$$\begin{aligned} a_{11} &= \frac{E_{\Delta,n,+}\mu_{n,-} + E_{\Delta,n,-}\mu_{n,+}}{\sqrt{E_{\Delta,n,+}^2 + \mu_{n,+}^2}\sqrt{E_{\Delta,n,-}^2 + \mu_{n,-}^2}}, & a_{12} &= \frac{E_{\Delta,n,+}\nu_{n,-} + E_{\Delta,n,-}\mu_{n,+}}{\sqrt{E_{\Delta,n,+}^2 + \mu_{n,+}^2}\sqrt{E_{\Delta,n,-}^2 + \nu_{n,-}^2}}, \\ a_{21} &= \frac{E_{\Delta,n,+}\mu_{n,-} + E_{\Delta,n,-}\nu_{n,+}}{\sqrt{E_{\Delta,n,+}^2 + \nu_{n,+}^2}\sqrt{E_{\Delta,n,-}^2 + \mu_{n,-}^2}}, & a_{22} &= \frac{E_{\Delta,n,+}\nu_{n,-} + E_{\Delta,n,-}\nu_{n,+}}{\sqrt{E_{\Delta,n,+}^2 + \nu_{n,+}^2}\sqrt{E_{\Delta,n,-}^2 + \nu_{n,-}^2}}. \end{aligned}$$

References

- [1] L. Allen and J. H. Eberly : Optical Resonance and Two-Level Atoms, Wiley, New York, 1975.
- [2] P. Meystre and M. Sargent III : Elements of Quantum Optics, Springer-Verlag, 1990.
- [3] Claude Cohen-Tannoudji, J. Dupont-Roc and G. Grynberg : Atom-Photon Interactions ; Basic Processes and Applications, Wiley, New York, 1998.
- [4] C. Montoe, D. Leibfried, B. E. King, D. M. Meekhof, W. M. Itano and D. J. Wineland : Simplified Quantum Logic with Trapped Ions, Physical Review A 55, R2489, quant-ph/9612042.
- [5] W. M. Itano, C. Montoe, D. M. Meekhof, D. Leibfried, B. E. King and D. J. Wineland : Quantum harmonic oscillator state synthesis and analysis, in Proc. Conf. on Atom Optics, San Jose, CA, Feb. 1997, edited by M. G. Prentiss and W. D. Phillips, quant-ph/9702038.
- [6] H. Moya-Cessa, A. Vidiella, J. A. Roversi, D. S. Freitas and S. M. Dutra : Long-time-scale revivals in ion traps, Phys. Rev. A 59(1999), 2518.
- [7] M. Frasca : Theory of dressed states in quantum optics, Phys. Rev. A 60(1999), 573, quant-ph/9811037.
- [8] M. Frasca : Rabi oscillations and macroscopic quantum superposition states, Phys. Rev. A 66(2002), 023810, quant-ph/0111134.
- [9] M. Frasca : A modern review of the two-level approximation, to appear in Annals of Physics, quant-ph/0209056.
- [10] M. Frasca : Perturbative results on localization for a driven two-level system, cond-mat/0303655.

- [11] Y. Nakamura, Yu. A. Pashkin and J. S. Tsai : Rabi Oscillations in a Josephson–Junction Charge Two–Level System, *Phys. Rev. Lett.*, 87(2001), 246601.
- [12] K. Fujii : Introduction to Grassmann Manifolds and Quantum Computation, *J. Applied Math*, 2(2002), 371, quant-ph/0103011.
- [13] K. Fujii : Mathematical Structure of Rabi Oscillations in the Strong Coupling Regime, *J. Phys. A* 36(2003), 2109, quant-ph/0203135.
- [14] K. Fujii : N–Level System Interacting with Single Radiation Mode and Multi Cat States of Schrödinger in the Strong Coupling Regime, quant-ph/0210166.
- [15] K. Fujii : A Generalized Hamiltonian Characterizing the Interaction of the Two–Level Atom and both the Single Radiation Mode and External Field, quant-ph/0303118.
- [16] K. Fujii : Cavity QED and Quantum Computation in the Strong Coupling Regime, quant-ph/0305155.
- [17] K. Fujii : Introduction to Coherent States and Quantum Information Theory, quant-ph/0112090.
- [18] K. Fujii : Two–Level System and Some Approximate Solutions in the Strong Coupling Regime, quant-ph/0301145.
- [19] K. Fujii : How To Treat An N–Level System : A Proposal, quant-ph/0302050.
- [20] K. Fujii, K. Higashida, R. Kato and Y. Wada : N Level System with RWA and Analytical Solutions Revisited, quant-ph/0307066.
- [21] J. C. A. Barata and W. F. Wreszinski : Strong Coupling Theory of Two Level Atoms in Periodic Fields, *Phys. Rev. Lett.* 84(2000), 2112, physics/9906029.

- [22] A. Santana, J. M. Gomez Llorente and V. Delgado : Semiclassical dressed states of two-level quantum systems driven by nonresonant and/or strong laser fields, J. Phys. B 34(2001), 2371, quant-ph/0011015.
- [23] C. E. Creffield : Location of crossing in the spectrum of a driven two-level system, Phys. Rev. B 67(2003), 165301. cond-mat/0301168.
- [24] K. Funahashi : Explicit Construction of Controlled-U and Unitary Transformation in Two-Qudit, quant-ph/0304078.
- [25] E. Solano, G. S. Agarwal and H. Walther : Strong-driving-assisted multipartite entanglement in cavity QED, Phys. Rev. Lett, 90(2003), 027903, quant-ph/0202071.